

POST-OPERATIVE INSTRUCTIONS

Total Hip Replacement

Hip Arthroplasty

Questions or a problem after surgery? Call Dr. Wichman's office at **414-479-7000**.

Diet

1. Begin with clear liquids and light foods (jello, soup, toast, etc...)
2. Progress slowly back to your pre-op diet as tolerated
3. Try to avoid taking narcotics on an empty stomach to avoid nausea

Wound Care

1. Take good care of your operative dressing.
2. Your dressing is water-tight and should remain in place until your first post-op visit.
3. You may shower after surgery.
4. To avoid infections, keep surgical incisions clean and dry. –NO immersion of leg in water (ie: bath or pool)

Medications

1. Pain medication is injected into the wound and hip joint during surgery. This will wear off in approximately 8-12 hours from surgery.
2. Most patients will require a period of narcotic pain medication. This should be taken as directed on the bottle
3. Norco (hydrocodone/tylenol) or Percocet (oxycodone/Tylenol)
4. Take 1-2 tablets every 4-6 hours as needed for pain.
5. Maximum of 12 pills per 24 hour period
6. Do NOT take additional Tylenol (acetaminophen) while taking these medications. Many products contain this medication (cough meds, cold meds) and can lead to liver damage.
7. Common side effects of narcotic pain medications are nausea, drowsiness, and constipation. Use a stool softener (Colace, senokot) or laxative (Miralax) as needed
8. You will be prescribed ONE of the following medications to thin your blood to reduce the risks of a blood clot/DVT. The choice of medications is based on your medical history and risk factors for blood clot. This will be determined by Dr. Wichman at the time of surgery
 - Aspirin (81mg or 325mg, as instructed by your doctor) twice daily
 - Lovenox 40mg subcutaneous injection ONCE daily
 - Eliquis 2.5mg orally twice daily
 - Other (per Dr. Wichman's discretion)
9. If you are having problems with nausea and vomiting, call the office to see if the medication should be changed or additional medications prescribed.
10. Journavx (suzetrigine): If you were prescribed Journavx, take 2 tablets (your one-time starting dose) with a small sip of water before leaving home the morning of surgery. Then take 1 tablet every 12 hours, starting 12 hours after your first dose, as directed. Take the starting dose on an empty stomach, and avoid grapefruit while taking this medication.
11. Do not drive a car or operate heavy machinery while taking narcotics.

Activity

1. Unless otherwise specified, you may bear weight as tolerated immediately after surgery.
2. No driving until instructed by medical team
3. Wear your compression stockings on BOTH legs until your post-op visit.
4. Your first post-operative visit will be with a PA, typically 7-14 days after surgery. Physical therapy will be arranged at that time.
5. It is critical that you follow the appropriate precautions after surgery
6. Posterior approach: For SIX (6) weeks- NO crossing your legs or bending past 90 degrees. Sleep with 1-2 pillows between your knees.
7. SuperPath- For SIX (6) weeks- NO flexion past 120 degrees.

Ice Therapy

1. Begin immediately after surgery
2. Use ice machine or ice packs every 2 hours for 20 minutes until your first post-operative visit.
3. If an ice machine was prescribed, may use continuously until first post-operative visit. Remember to protect the skin to avoid thermal injury (frostbite)

Exercise

1. Begin exercises on post-operative day one. (Straight leg raises, quad sets, heel slides and ankle pumps)
2. REMEMBER, there will be some pain associated with your exercises, utilize your pain medications to allow you to maximize your motion.
3. Complete exercises 3-4 times daily until your first post-operative visit.
4. Perform ankle pumps throughout the day to reduce risk of blood clot in your calf.

When to Call Us

Contact Dr. Wichman at 414-479-7000 if any of the following are present:

- Painful swelling or numbness
- Unrelenting pain
- Redness around incisions
- Fever (above 101.5°F) - It is not uncommon to have a low-grade fever for the first day or two following surgery.
- Color changes in foot, ankle, or lower extremity
- Continuous drainage or bleeding from incisions (a small amount of drainage is expected.)
- Difficulty breathing
- Excessive nausea or vomiting

If you have an emergency after hours or over the weekend, call 414-479-7000 to be connected to the "on-call" physician or physician assistant. Do NOT call the hospital or surgery center.

If you have an emergency that requires immediate attention, call 911 or proceed to the nearest Emergency Room.